

FORAGE CROPS

J. Soroka and J. Otani

ALFALFA BLOTCH LEAFMINER *Agromyza frontella* (Rond.)

Economic Thresholds

Alfalfa hay crops
- 40% of leaflets show pinhole feeding (OMAFRA)

Alfalfa seed crops
- unknown

Biological Control -

A species of parasitic wasp, *Dacnusa dryas* (Nixon), has kept the leafminer in check in eastern Canada in most localities and years. Chemical control of the pest may decimate populations of this biocontrol agent.

Chemical Control –

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	References
dimethoate Cygon 480EC, AG Lagon 480	264	0.55 L	0.223 L	2	-
malathion Fyfanon 50% Malathion 85E	1375 1143	2.75 L 1.345 L	1.11 L 0.544 L	7 7	-
phosmet Imidan 50 WP Imidan 70 WP	1125 1120	2.25 kg 1.60 kg	0.91 kg 0.65 kg	7 7	-

Note: Insecticides should be applied no later than the pinhole stage of feeding, usually in mid- to late May.

Restrictions -

- crop in bloom Do not apply during bloom. See Section in Guide regarding hazard to bees.
- dimethoate Use a minimum water volume of 200 L/ha with ground equipment. Do not apply more than once per season. Ground and aerial application is permitted.
- malathion Greatest efficacy at air temperatures above 20°C. Use a minimum water volume of 275 L/ha. Repeat application may be necessary. Some products are restricted to ground application only; others may be applied by ground or air.

phosmet Use water volumes of 200-500 L/ha. Apply when first signs of infestation are visible. Do not apply more than once per cutting. Do not make more than three applications per season. Do not apply by air. Restricted entry interval is 5 days.

ALFALFA CATERPILLAR *Colias eurytheme* (Bois.)

Various lepidopteran larvae feed on legume and grass forages, usually with little impact because populations are kept low by natural biological controls. Normal forage harvesting disrupts larval life cycles by preventing many larvae from maturing. Alfalfa caterpillar occurs fairly frequently in alfalfa on the prairies but has not been recorded as causing economic harm (1).

Economic thresholds – unknown

Chemical control -

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
carbaryl Sevin XLR	963-2247	2.5-5.25 kg	1.01-2.12 kg	0	-

Restrictions -

carbaryl Remove cattle from area while spraying. Re-introduce beef cattle after 1 day, dairy cattle 2 days. Repeat in 7 to 14 days if necessary. Ground or aerial application.

Reference -

1. Beirne, Mem. Entomol. Soc Can. No. 78, 1971.

ALFALFA LOOPER *Autographa californica* (Speyer)

Economic Thresholds

No economic thresholds have been established for alfalfa.

Biological Control -

The alfalfa looper is usually held in check by its natural parasites and predators, and by viral diseases that kill the larvae before they can pupate.

Cultural Control -

Early cutting may give satisfactory control because larvae cannot survive on dried alfalfa forage. Alfalfa should be inspected within 2 to 3 days after cutting to determine if loopers are feeding on regrowth. Insecticide application in alfalfa is rarely warranted.

Chemical Control –

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
naled Dibrom 480 EC	950-1900	1.1-2.2 L	0.445-0.890 L	4	-

Restrictions

crop in bloom Do not apply to crops in bloom. See Section in Guide regarding hazard to bees.

naled Ground or aerial application. For ground application, dilute with water to 100-300 L/ha unless otherwise stated. For aerial application, dilute with water to 10-30 L/ha unless otherwise stated. When using maximum rate, the area treated is limited to 200 ha/day. Re-entry period is 48 hr. Do not apply when temperatures exceed 32°C.

ALFALFA WEEVIL *Hypera postica* (Gyll.)

Economic Thresholds

Alfalfa hay crops (6-9)
 - 25-50% of leaves on the upper one-third of the stem show damage (6)
 - 50-70% of terminals show injury (9)

Alfalfa seed crops (10)
 - 20-25 larvae/sweep (90°=straight sweep)
 - 35-50% of foliage tips show damage

Biological Control -

The parasitic wasp parasitic wasp *Bathyplectes curculionis* (Thomson) is now found in Alberta and Saskatchewan. Chemical control of the pest may decimate populations of this biocontrol agent.

Cultural Control -

Cut the first hay crop early. If damage reappears in new growth, insecticide application may be necessary.

Chemical Control -

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	References
carbaryl Sevin 50W	1625	3.25 kg	1.315 kg	0	-
chlorantranilprole Coragen	75-100	0.375 -0.50 L	0.152-0.202 L	0	-
deltamethrin Decis 5 EC	10-12.5	0.2-0.25 L	0.08-0.10 L	20	11, 12
dimethoate	204	0.425 L	0.172 L	2	-

Cygon 480EC, 480AG, Lagon					
λ-cyhalothrin Matador 120EC Silencer	10	0.083 L	0.034 L	3	-
malathion Malathion 500, 500E Malathion 85E Malathion ULV	1000-1500 935-1143	2-3 L 1.1-1.345 L 1.1	0.809-1.214 L 0.445-0.544 L 0.445	7 7 5	1, 2
phosmet Imidan 50 WP Imidan 70 WP	1125 1120	2.25 kg 1.6 kg	0.911 kg 0.648 kg	7	-

Forage Crop If application is made to heavy growth, 110-220 L water/ha may be required for adequate coverage.

Seed Crop Chlorantraniliprole, deltamethrin, and λ- cyhalothrin are less toxic to pollinators than other products. Keep leafcutting bees off the crop until the cyhalothrin-λ dries completely or at least 24 hours following deltamethrin application.

Restrictions -

crop in bloom Do not apply insecticide when bees are pollinating. See Section in Guide regarding hazard to bees.

carbaryl Remove cattle from area when spraying. A repeat application may be made in 7 to 14 days if necessary. If pre-treatment damage is extensive, cut and make application to stubble. Ground application only.

chlorantraniliprole Suppression only. Do not make more than one application per cutting. Recommended to wait one day between application and feeding to livestock. Not registered specifically for alfalfa seed production. Ground application only.

deltamethrin For use on seed crops only. Avoid application when daytime temperatures are greater than 25°C. One application per year. Apply by ground only.

dimethoate For reduction of alfalfa weevil larvae. Do not apply when alfalfa is in bloom. Remove cattle before spraying. Maximum 2 applications per season. Ground or aerial application is permitted.

λ-cyhalothrin Do not apply within 3 days of grazing; on unimproved pastures the recommended pre-grazing interval is 7 days. Allow a 7 day interval between treatments. Do not use more than 3 applications per season. Alfalfa seed from treated crops must not be used for production of alfalfa sprouts. Ground or aerial application of Matador is permitted. Do not make more than 1 application of 83 mL/ha of the allowed seasonal total by air. Ground application only for Silencer.

- malathion Some products list control of alfalfa weevil larvae only. Greatest efficacy at air temperatures above 20°C. Apply when 75% of leaves show feeding damage. Do not apply when alfalfa is in bloom. Ground or aerial application.
- phosmet Apply in 200-500L water per ha. Do not apply more than once per cutting. Do not make more than 3 applications per season. Restricted entry interval is 5 days. Consult local agricultural authorities regarding proper time of spray application. Ground application only.

References -

1. Bass and Knapp, J. Econ. Entomol. 59:648, 1966.
2. Swailes and McDonald, Pest. Res. Rep. 1964:177.
3. Ellis, Pest. Res. Rep. 1976:167.
4. Richards and Charnetski, Pest. Res. Rep. 1976:168.
5. Harper, Can. Entomol. 110:891, 1978.
6. Bereza, Ontario Agdex 121/622. 4 pp., 1977.
7. Hastings and Pepper, J. Econ. Entomol. 46:785, 1953.
8. Hintz et al., J. Econ. Entomol. 69:749, 1976.
9. Carpenter, J. Econ. Entomol. 63: 1602, 1970
10. Johansen, Pacific North West Ext. Publ. No. 128, 1980.
11. Charnetski and Schaber, Pest. Res. Rep. 1980:160.
12. Charnetski, Pest. Res. Rep. 1983:154.

ARMYWORMS including true armyworm *Mythimna unipuncta*, Hay, fall armyworm *Spodoptera frugiperda* (J.E. Smith) and others

Armyworms can cause severe damage to cereal and grass crops or pastures in outbreak situations, but such years are often interspersed with several years of little damage. Infestations can be quite localized. True armyworm moths are annual migrants to the prairies from the United States.

Economic thresholds – are unknown in grasses

Biological control - Many species of parasitoids and predators, as well as bacterial and fungal diseases and even birds feed on armyworms. The control of such natural enemies may come too late however, to reduce economic damage in high infestations (1)

Chemical control -

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
carbaryl Sevin XLR	963-2247	2.5-5.25 kg	1.01-2.12 kg	0	-
chlorantraniliprole Coragen	50 - 75	0.25-0.375 L	0.101-0.152 L	0	

Restrictions -

carbaryl Remove cattle from area while spraying. Re-introduce beef cattle after 1 day, dairy cattle 2 days. Repeat in 7 to 14 days if necessary. Ground or aerial application.

chlorantraniliprole For grass seed production only. Crop residue is not to be fed to livestock. Do not apply more than four times per year or more than once every 7 days, and do not exceed application of more than 1.125L/ha per season.

Reference -

1. Beirne, Mem. Entomol. Soc Can. No. 78, 1971.

BEET WEBWORM <i>Loxostege sticticalis</i> (L.) and other webworms
--

Beet webworm has a wide host range, often preferring to feed on weeds in a crop than the crops itself. It is a sporadic pest that causes damage only in some years in some locations. Normal forage harvesting disrupts life cycles of Lepidoptera such as beet webworm by preventing many larvae from maturing.

Economic thresholds – unknown

Chemical control -

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
carbaryl Sevin XLR	963-2247	2.5-5.25 kg	1.01-2.12 kg	0	-

Restrictions -

carbaryl Remove cattle from area while spraying. Re-introduce beef cattle after 1 day, dairy cattle 2 days. Repeat in 7 to 14 days if necessary. Ground or aerial application.

BLISTER BEETLE <i>Epicauta</i> spp., <i>Lytta</i> spp.

Damage, although usually localized, can occur quickly and mainly in dry situations when grasshoppers are present. Blister beetle larvae eat grasshopper eggs.

Chemical control-

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
carbaryl Sevin 50W	1125-1625	2.25-3.25	0.91-1.32 kg	0	-
Sevin XLR	1070-1712	2.5-4.0	1.01-1.62 kg	2	

Restrictions -

carbaryl

Remove cattle from area while spraying. Re-introduce beef cattle after 1 day, dairy cattle 2 days. Repeat in 7 to 14 days if necessary. Ground application only of Sevin 50W, ground or aerial application of Sevin XLR.

BROMEGRASS SEED MIDGE <i>Contarinia bromicola</i> (Marik & Agaf.)
--

Cultural Control -

Thorough burning of stubble and trash in early spring or before regrowth in late summer provides some control. (1-2) Parasitism by a species of *Tetrastichus* (Eulophidae) can vary from 30% to 75% (3).

References -

1. Knowles, Forage Notes 18(2), 1973; 19(2), 1974.
2. Neuman and Manglitz, Agric. Expt. Sta., Univ. Neb., Res. Bull. 252, 1972.
3. Curry, Knowles and Waddington, Can. Entomol. 115:(1) 75-79, 1983.

ESSEX (EUROPEAN) SKIPPER <i>Thymelicus lineola</i> (Ochs.)

on timothy

Chemical control-

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
<i>Bacillus thuringiensis</i> Dipel WP, 2X-DF, BioProtec 3P BioProtec CAF	80-176	0.275-0.550 kg	0.111-0.223 kg		-
		0.55-0.70 L	0.223-0.283 L	1	

Restrictions -

Bacillus thuringiensis Apply at first sign of infestation and repeat at 7 to 10 day intervals when needed to maintain control. May be applied up to the day of harvest. Use in a minimum of 300 litres of water per hectare. Use diluted spray mixtures within 12 hr. Ground or air application.

GRASSHOPPERS Orthoptera:Acrididae
--

on alfalfa, clover, forage grasses, pastures and rangelands

Economic Thresholds - See Cereal Crops Section. Unless the forage crops are in their establishment year, economic thresholds should be closer to ETs for roadsides than those for field crops.

Chemical Control -

Active ingredient Product	Crop	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Refer- ence
carbaryl Sevin XLR	aerial or ground: alfalfa, clover	535-1070	1.25-2.50	0.505-1.01 kg	2	-
	ground: forage grasses, pastures, rangelands	514-984	1.2-2.3 kg for nymphs or sparse vegetation	0.486-0.931 kg	1 (beef cattle) – 2 (dairy cattle)	-
		984-1498	2.3-3.5 kg for adults or dense vegetation	0.931-1.416 kg		
Sevin 50 WP	alfalfa, clover	550-1125	1.1-2.25 kg	0.445-0.910 kg	0	
carbaryl bran bait Sevin Bran Bait Eco Bran Bait	alfalfa, clover	40-100	2.0 kg (5%)	0.809 kg	2	-
	grass and legume forages, rangeland, pastures		2.0 of 2% carbaryl	0.809 - 1.161 kg	1 (beef cattle) – 2 (dairy cattle, harvest of forage crops)	
deltamethrin Decis 5 EC, F	rangeland, pastures	5.0 - 7.5 ground	0.08-0.15 L in 100-200 L water	0.032-0.061 L	1	
		7.5 aerial	0.12-0.15 L in 11- 22 L water	0.0[49-0.061 L		
dimethoate Cygon 480EC, 480 AG Lagon 480 E	alfalfa seed and forage, forage crops (grasses, hay, clover), pastures	204-480	0.425-0.55 L – low infestations	0.172 - 0.222 L – low infestations	2 (low rates) – 28 (high rate)	1-4
			0.550 L -nymphs	0.223 L nymphs		
			0.850 – 1.0 L - adults	0.344- 0.40 L adults		
λ -cyhalothrin Matador, Silencer, Warrior	alfalfa/grass, timothy for seed or hay, pasture	7.6 - 10 ground 10 aerial	0.063-0.083 L (ground)	0.025-0.034 L (ground)	3-14	-
			0.083 L (aerial)	0.034 L (aerial)		

malathion Malathion 50 EC, 500 E	alfalfa, clovers, grasses or legumes for hay	850-1250	1.7-2.5 L	0.69-1.01 L	7	-
ULV	grass in pasture, rangeland	530-830	0.55-0.85 L	0.22-0.34 L	0	-
Malathion 85E	alfalfa	935 - 1143	1.10- 1.345 kg	0.44 – 0.55 kg	7	-
naled Dibrom	rangeland, pastures	475 - 735 young GH 605 - 865 adult GH	0.55 - 0.85 L 0.7 - 1.0 L	0.22 -0.34 L 0.28 - 0.40 L	0	-
<i>Nosema locustae</i> Nolo Bran Bait	crop and rangeland		minimum 1.12 kg	minimum 0.453 kg	0	-

Note: Use lower rates on younger grasshoppers. Best control is achieved when product is applied to young grasshoppers. If insect pressure is high spraying a 15 m strip around the field may aid in control.

Restrictions -

- crop in bloom Do not apply to crops in bloom. See Section in Guide regarding hazard to bees.
- carbaryl Remove cattle from area while spraying. Dilutions of higher than 1:11 are not recommended when residual insect control from the wash-off is desired. Sevin XLR - ground or aerial application, Sevin WP – ground application only.
- carbaryl
bran bait Apply by ground only. Rain shortly after application will decrease efficacy of product. Eco Bran Bait - repeat applications only as necessary but not more frequently than once a week during periods of heavy infestation and once every two weeks during moderate to low infestation. Presence of product on flowering crops such as alfalfa and clover will not harm foraging honey or leafcutter bees.
- cyhalothrin-λ Do not use more than 3 applications per year. Air or ground application, but do not make more than 1 application by air per year. Do not use alfalfa seed from treated crops for production of alfalfa sprouts. Allow 7 days between applications. Timothy for seed or hay – apply by ground only. Use sufficient water for thorough coverage, 100 to 200 L/ha by ground sprayer. Treated crops may be fed to non-lactating dairy animals and other livestock after a 3 day interval from application to harvest or foraging; **exception** – do not apply within 14 days of timothy harvest. Apply in 100 to 200 L of water per hectare.
- deltamethrin Do not make more than three applications per year, with a maximum of two aerial applications. Use higher rates for older insects and/or dense crop canopy.

dimethoate	Do not apply during bloom. When spraying honey-producing crops (alfalfa, red clover, and alsike) spray at least 5 days before bloom appears and do not introduce hives until full bloom. Follow provincial forecast. Remove cattle before spraying. Ground or aerial application.
malathion	Greatest efficacy at air temperatures above 20°C. Pasture and rangelands treated with malathion may be grazed or harvested on day of treatment. No application within 7 days of harvest or pasturing of alfalfa, clovers. Do not apply to alfalfa in bloom. Apply when 75% of foliage shows feeding damage. Repeat as necessary. Ground or aerial application.
naled	Use sufficient water to provide thorough coverage and contact spray. Animals may be present during treatment. Do not apply when temperatures exceed 32°C. Ground or aerial application.

Nosema lacustae Grasshopper death will begin in 3-6 weeks; the pathogen will remain in the grasshopper population for several years. Ground or aerial application.

References -

1. Holmes et al., J. Econ. Entomol. 58:77, 1965.
2. Banham, Pest. Res. Rep. 1964:134.
3. McDonald and McKinlay, Pest. Res. Rep. 1964:132.
4. Dolinski et al., Pest. Res. Rep. 1973:185.
5. Mukerji et al., Can. Entomol. 1981:707.
6. Johnson et al., Pest. Res. Rep. 1985: 174.

LEAFHOPPERS (Cicadellidae), including POTATO LEAFHOPPER *Empoasca fabae* (Harris)

Many species of leafhopper are present in prairie forage crops but damage either by direct feeding or by disease transmission is rare. However, in some areas of the eastern North America leaf hoppers are rated among the most serious pests of alfalfa. Potato leafhopper damage symptoms appear as stunting as well as yellowing of the leaves in a v-shaped pattern starting at the tip of a leaflet. Damage is most severe in new seedlings and in regrowth in hot, dry weather.

Economic thresholds – vary with commodity prices and application costs. Historic thresholds are: if alfalfa is <9 cm in height, the economic threshold for potato leafhoppers is 0.2 adults/sweep, <15 cm=0.5 adult PLH, <25 cm= 1.0 adult PLH, and <36 cm=2.0 adult PLH/sweep.(1)

Cultural Control –

Cut the first hay crop early. Some alfalfa cultivars have resistance to leafhoppers.

Chemical Control –

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
------------------------------	-------------------	----------	------------	----------------------------------	-----------

carbaryl Sevin 50 WP Sevin XLR	1012-1712	2.25-3.25 kg 2.5-4.0 L	0.91-1.32 kg 1.01-1.612 L	0 2	-
dimethoate Cygon, Lagon	204	0.425 L	0.172 L	2	-
cyhalothrin-λ Matador, Silencer	10	0.083 L	0.034 L	3	-
malathion Malathion 500 EC Malathion 500 Malathion 85E	935 – 1500	2-3 L 1.1 – 1.345 L	0.809-1.214 L 0.445-0.554 L	7	-
naled Dibrom	950-1900	1.1-2.2 L	0.445-0.890 L	4	-

Restrictions

- crop in bloom Do not apply to crops in bloom. See Section in Guide regarding hazard to bees.
- carbaryl Remove cattle from area when spraying. Reintroduce beef cattle after 1 day, dairy cattle after 2 days. Ground or aerial application.
- dimethoate Remove cattle before spraying. Use sufficient water for good coverage. Repeat only if necessary. Ground or aerial application.
- λ-cyhalothrin Do not apply within 3 days of grazing; on unimproved pastures the recommended pre-grazing interval is 7 days. Allow a 7 day interval between treatments. Do not use more than 3 applications per season. Alfalfa seed from treated crops must not be used for production of alfalfa sprouts. Matador -ground or aerial application; Silencer – ground application only. Do not make more than 1 application of 83 mL/ha of the allowed seasonal total by air.
- malathion Greatest efficacy at air temperatures above 20°C. Apply when 75% of foliage shows feeding damage. Repeat as necessary. Ground or aerial application.
- naled Ground or aerial application. For ground application, dilute with water to 100-300 L/ha unless otherwise stated. For aerial application, dilute with water to 10-30 L/ha unless otherwise stated. When using maximum rate, the area treated is limited to 200 ha/day. Re-entry period is 48 hr. Do not apply when temperatures exceed 32°C.

Reference -

1. Metcalf and Luckmann. 1994. Introduction to Integrated Pest Management. pp 469-504.

LESSER CLOVER LEAF WEEVIL *Hypera nigrirostris* (F.)

Economic thresholds

Red clover

- third and fourth instar larval densities greater than 3 per five shoots could damage up to 50% of buds and flower heads (1)

Chemical control - in established red clover fields for seed production only

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
deltamethrin Decis 5EC	12.5	0.250 L in 100-300 L	0.101 L	n/a	-

Restrictions

crop in bloom Do not apply insecticide when bees are pollinating. See Section in Guide regarding hazard to bees.

deltamethrin For suppression. Do not make more than two applications per year. Apply two times at a 14 day interval, at early vegetative and when crop is budding. Do not use treated crop for feed or forage. Restricted entry interval 12 hours. Ground application only.

Reference -

1. Weiss and Gillott, Can. Entomol. 125:831-837, 1993.

PEA APHID *Acyrtosiphon pisum* Harr. and other aphids

Economic Thresholds -

1430 aphids/sweep did not reduce the forage yield of irrigated alfalfa. (2)

Caged alfalfa initially infested with 100-200 aphids/plant produced less forage and had lower carotene contents than uninfested plants. (4)

100 - 200 aphids/ 180° sweep when seed crop is drought-stressed, plants are wilting, and seed is still developing (before mid-August)

Cultural Control -

On irrigated alfalfa, control may be unnecessary if adequate water is provided to the plants. (2)

Biological Control -

High winds, heavy rain and hail can cause pea aphid numbers to decline drastically. Predators such as ladybird beetles and others, parasitic wasps, and entomopathogenic diseases often cause aphid population crashes to occur, especially late in the season.

Chemical Control -

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
dimethoate	204	0.425 L	0.172 L	2	1, 3

Cygon, Lagon					
λ-cyhalothrin Matador, Silencer	10	0.083 L	0.034 L	3	-
flonicamid Beleaf 50 SG	0.06-0.08	0.12-0.16	0.05-0.06	-	-
malathion Malathion 500, 500E	895 – 1500	2 -3 L	0.809-1.214 L	7	-
Malathion 85E		1.1 – 1.345 L	0.445-0.544 L		
naled Dibrom	950-1900	1.1-2.2 L	0.445-0.890 L	4	-

Note: Ensure thorough coverage. Treat if population is large enough to stunt plants or if heavily infested hay is to be dehydrated.

Restrictions -

Do not apply when pollinators are present. See Section in Guide regarding hazard to bees.

- dimethoate Remove cattle before spraying. Maximum number of applications per season - two. Suppression only of Russian what aphid. Ground or aerial application.
- flonicamid For alfalfa seed production only. Maximum of three applications per season; minimum of 7 day interval between applications. Do not use alfalfa seed from treated crops for human consumption
- λ-cyhalothrin Maximum of three applications per season; minimum of 7 day interval between applications. Do not use alfalfa seed from treated crops for human consumption.. Air or ground application of Matador, but do not make more than 1 application by air per year. Silencer - ground application only.
- malathion Do not apply to alfalfa in bloom. Remove cattle before spraying. Greatest efficacy at air temperatures above 20°C. Application by ground or air.
- naled Ground or aerial application. For ground application, dilute with water to 100-300 L/ha unless otherwise stated. For aerial application, dilute with water to 10-30 L/ha unless otherwise stated. When using maximum rate, the area treated is limited to 200 ha/day. Maximum two applications per season. Re-entry period is 48 hr. Do not apply when temperatures exceed 32°C.

References -

1. Harper, Can. Entomol. 110:891, 1978.
2. Hobbs et al., Can. Entomol. 93:801, 1961.
3. McDonald and Harper, Can. Entomol. 110:213, 1978.
4. Harper and Lilly, J. Econ. Entomol. 59:1426, 1966.

PLANT BUGS IN ALFALFA SEED FIELDS

Including Lygus bugs, Alfalfa plant bug, Superb plant bug, and Plagiognathus bug.

Economic Thresholds -

8 lygus bugs/sweep or 4 alfalfa plant bugs/sweep or 5 nymphs/sweep of any or all species of plant bugs, when the alfalfa is in bud or in bloom. (7-9)

Chemical application should be timed so that the majority of nymphs are third instars or older.

Cultural Control -

Burning of alfalfa stubble and debris in early spring controls all species except lygus bugs. (3, 5, 15, 16)

Chemical Control -

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	References
acetamiprid Assail 70 WP	60-120	0.085-0.170 kg	0.0352-0.0688 kg	1	-
deltamethrin Decis 5EC	10 - 12.5	0.20-0.25 L	0.081-0.101 L	20	10-14
dimethoate – forage – Cygon 480, Lagon seed - Cygon 480, Lagon	204 – 528	0.425 L 1.1 L	0.172 L 0.445 L	2 28	1, 6
λ-cyhalothrin Matador, Silencer	10	0.083 L	0.034 L	3	-
malathion Malathion 500 Malathion 85E	935 – 1500	2.0-3.0 L 1.10-1.345 L	0.809-1.214 L 0.445-0.544 L	7	-
naled Dibrom	950 - 1900	1.1-2.2 L	0.445-0.890 L	4	-
novaluron Rimon	90	0.835 L	0.338 L	14	-

Note: For control of lygus bugs in burned fields, apply dimethoate or deltamethrin just before alfalfa begins to bloom.

When putting out or replacing leafcutting bees on the crop, wait 7 days after applying dimethoate and wait 1 day after applying deltamethrin.

In unburned fields for control of all plant bugs, apply dimethoate no later than the beginning of flower bud formation.

If the alfalfa is in bloom and bees are in the crop, apply cyhalothrin- λ in late evening when bees are not foraging.

Restrictions –

acetamiprid	Seed production only. Control of alfalfa plant bug, suppression of Lygus bug. Apply prior to bloom up to the time when 50% of seed pods are ripe. Begin when adults and/or 4 th -5 th instar nymphs have reached economic thresholds. Use the higher rate for heavier infestations. Maximum 3 applications per season. Do not apply more than once every 7 days. Do not cut treated fields for hay/forage or graze treated fields. Do not exceed a total of 357 g active ingredient (510 g product) per ha per season. Ground application only.
deltamethrin	Seed production only. Do not feed treated forage within 90 days of harvest. Do not apply more than once annually. Use higher rates when alfalfa weevils are present. Not recommended for alfalfa plant bug control. Apply in 100-300 L/ha water. Ground application only.
dimethoate	Do not graze or harvest for forage within 2 days following treatment. Ground or aerial application. Allow at least 7 days before placing leaf cutting bees in the field. Do not apply more than once per season.
λ -cyhalothrin	Do not graze or feed treated forage or pasture to lactating dairy cows. Allow 14 days between application and cutting for forage or grazing of non-lactating dairy animals or other livestock. Do not use more than 3 applications per year. Do not use alfalfa seed from treated crops for production of alfalfa sprouts. Aerial and ground (Matador) or ground (Silencer) application.
malathion	Greatest efficacy at air temperatures above 20°C. Apply when 75% of foliage shows damage. Do not apply to alfalfa in bloom. Ground or aerial application permitted.
naled	Ground or aerial application. For ground application, dilute with water to 100-300 L/ha unless otherwise stated. For aerial application, dilute with water to 10-30 L/ha unless otherwise stated. When using maximum rate, the area treated is limited to 200 ha/day. Re-entry period is 48 hr. Do not apply when temperatures exceed 32°C.
novaluron	For alfalfa seed production only. For control of plant bug nymphs only. Avoid application to crops in bloom if bees are in the area. Use only on pure stands of alfalfa to be harvested for seed. Do not allow livestock to graze treated fields. Do not use treated crops or crop residue for livestock feed. Alfalfa seed from treated fields may not be used for growing sprouts for human or animal consumption. Ground application only.

References -

1. Craig, Pest. Res. Rep. 1971:167.
2. Kolach and Senkow, Pest. Res. Rep. 1972:151.
3. Lilly and Hobbs, Can. J. Plant Sci. 42:53, 1962.
4. Craig, Pest. Res. Rep. 1973:168.
5. Craig, Agric. Can. Publ. 1935, 1973.

6. Harper, Can. Entomol. 110:891, 1978.
7. Craig and Lasiuk, Pest. Res. Rep. 1971:167.
8. McMahon, Rep. B.C. Agron. Assoc. 1950:58.
9. Sorenson, Utah Agric. Expt. Sta. Bull. 284,1939.
10. Charnetski and Schaber, Pest. Res. Rep. 1980:161.
11. Charnetski, Pest. Res. Rep. 1983:154.
12. Butts, Pest. Res. Rep. 1981:149; 1983:156.
13. Butts and Lippert, Pest. Res. Rep. 1982:166.
14. Cattellier and Wise, Pest. Res. Rep. 1982:167.
15. Schaber and Entz, J. Econ. Entomol. 81:668-672, 1987.
16. Craig, Pest. Res. Rep. 1980:163.

PLANT BUGS IN GRASS FIELDS

Plant bugs such as *Capsus cinctus* Kolenaty, *Trigonotylus coelestialis* (Kirk), others (See Silvertop of Grasses below)

Chemical Control -

Active Ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
dimethoate Cygon 480, Lagon	204	0.425 L	0.172 L	2	1

Restrictions -

dimethoate Do not graze, forage, or harvest 2 days after treatment. Ground or aerial application.

References -

1. Okuda, Pest. Res. Rep. 1988 p.134.

SILVERTOP OF GRASSES

Diverse causal agents. Insects associated with silvertop include grass plant bugs (e.g., *Capsus* species, *Trigonotylus* spp., *Irbisia* spp., *Labops* spp., others), thrips such as *Anaphothrips* species, and grain mites such as *Siteroptes graminum*. Diseases such as *Fusarium* have also been implicated.

Economic Thresholds -

Thresholds will vary according to seed prices, which fluctuate wildly. In general, grass fields exhibiting 10% or more silvertop should have remedial action taken the following year. Silvertop, once evident, cannot be treated or reversed in the current season.

Cultural Control -

Post-harvest burning or burning in the spring prior to new growth will destroy many plant bug eggs and adults. Do not burn creeping red fescue fields in the Peace River region because of potential damage to the subsequent seed crop. In this region, burn only after the last seed harvest and prior to entering a rejuvenation cycle. Low-mowing (below 3 cm) and removal of straw following harvest also decreases the incidence of silvertop.

Chemical Control -

At present, certain formulations of dimethoate are registered for control of plant bugs in forage crops (See Plant Bugs in Grass Fields above).

References -

1. Peterson and Vea, J. Econ. Entomol. 64:247-251, 1971
2. Kamm and Fuxa, J. Range Management 30:385-387, 1977
3. Okuda, Pest. Res. Rep. 1988:134.

SEED CHALCIDS including alfalfa seed chalcid *Bruchophagus roddi* (Guss.), clover seed chalcid *B. gibbis* (Boh.), trefoil seed chalcid *B. platpterus* (Wlkr), and sainfoin seed chalcid *Eurytoma brychidi* (Wlkr.)

Red clover thrips (red clover seed fields)

Economic Thresholds -

Damage is insignificant until 50-80 thrips/raceme are present. (1) Threshold levels occur only during years of early spring drought on dryland.

Reference -

1. Craig, unpublished data 1987, Agriculture and Agri-Food Canada, Saskatoon.

Seed chalcids do not affect legume forage production, but can cause up to 50% reduction in seed yield. Dryland crops suffer worse damage than irrigated fields. Likewise, damage is worse following a hot, dry summer.

Shallow cultivation of seed fields in the fall or early spring can bury many chalcid-infested seeds. Combine tailings and screenings should not be left in the field. Sow certified, chalcid-free seed.

References -

1. Soroka and Spurr, Can. Entomol. 130: 1-11, 1998.

SPIDER MITES *Tetranychus urticae* Koch and others

Spider mite populations can increase rapidly and cause damage in hot, dry conditions, usually towards the end of summer.

Economic Thresholds -

Unknown for Canadian prairies. In clover seed crops in Idaho, mite control is recommended in June or July when 25% of the leaves show feeding injury. (1)

Biological Control -

Mite populations are usually kept in check by low temperatures and rain, fungal diseases and predators such as ladybird beetles, predacious thrips and especially predacious mites.

Chemical Control – in alfalfa grown for seed

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
malathion Malathion 500E Malathion 85E	935-1375	2.25-2.75 L 1.1-1.345 L	0.910-1.13L 0.445-0.544 L	7	-
spiromesifen Oberon	120-240	0.5-1.0 L	0.202-0.405 L	n/a	-

Restrictions -

malathion Greatest efficacy at air temperatures above 20°C. Do not apply to alfalfa in bloom. Apply when 75% of foliage shows feeding damage. Repeat as necessary. Some products are restricted to ground application only; others may be applied by air.

spiromesifen Three applications may be required for control. Treat at minimum 7 day intervals. Do not apply when pollinators are actively foraging in the treatment area; apply before pollinators are placed in a field or after they have been removed. Ground and aerial application is permitted. Maximum allowed per crop season: 3 L/ha. Minimum application volume: 100 L/ha - ground; 50L/ha – aerial application.

References -

1. Baird, Homan and Bolz. 2002.

SPITTLE BUGS *Philaenus spumaris* (L.)

Spittle bugs are only an occasional pest of alfalfa, and not normally a cause for concern. Adults do not damage alfalfa. When nymphs hatch they form a spittle mass which is used to prevent desiccation and to protect from predators.

Economic Thresholds -

Unknown for Canadian prairies. In Wisconsin, an average of 1 spittle bug nymph/stem is necessary before control should be considered. (1)

Chemical Control -

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
malathion Malathion 500E Malathion 85E	935-1375	2.25-2.75 L 1.1-1.345 L	0.910-1.13L 0.445-0.544 L	7	-
endosulfan Thiodan 4 EC	300	0.75 L	0.303 L	30	-

Restrictions –

- malathion - Greatest efficacy at air temperatures above 20°C. Do not apply to alfalfa in bloom. Apply when 75% of foliage shows feeding damage. Repeat as necessary. Some products are restricted to ground application only; others may be applied by air.
- endosulfan Do not apply when bees are foraging. Apply soon after eggs hatch. Do not feed forage within 30 days of application. Ground application only.

References -

1. Baird, corn.agronomy.wisc.edu/Management/pdfs/IPMManual_2_Alfalfa.pdf page 10

SWEETCLOVER WEEVIL *Sitona cylindricollis* Fahr.

Economic Thresholds (7) -

In seedling sweet clover crops:

- 1 weevil adult/5 seedlings in cotyledon stage under slow growth conditions.
- 1 weevil adult/3 seedlings in cotyledon stage under normal growth conditions.

In newly-emerging 2nd year sweet clover:

- 9-12 weevil adults/plant.

Cultural Control -

Seedling stands:

- Locate new seedlings as far as possible from 2nd-year clover.

First-year stands in late summer:

- Defoliation of 1st-year clover by close-cutting, grazing, or weevil feeding during the critical period of mid-August to mid-September will severely reduce 2nd-year yield.
- Swath companion grain crops high.

Second-year stands:

- Second year clover will usually outgrow weevil damage; insecticide application is rarely necessary. Sweet clover silage and hay fields should be cultivated as soon as possible after the crop is taken; this kills the new-generation weevil larvae in the soil. (6,7)

Chemical Control -

In sweet clover seedling crops apply insecticide when very young seedlings are noticeably damaged by weevil feeding. In first-year clover in late summer and fall, apply insecticide to crop margins where weevils are concentrated.

Active ingredient Product	Rate (g AI/ha)	Rate /ha	Rate /acre	Preharvest Interval (days)	Reference
carbaryl Sevin 50W	1125-2000	2.25-4.0 kg	0.910-1.619 kg	0	-
dimethoate Cygon, Lagon	408 - 528	0.850-1.1 L	0.344-0.445 L	28	1-5
malathion Malathion 500, 500E Malathion 85E	625 - 1250	1.4-2.5 L 0.735-1.22 L	0.567-1.012 L 0.297-0.494 L	7	-

Restrictions –

carbaryl	Remove cattle from area when spraying. A repeat application may be made in 7 to 14 days if necessary. Ground or aerial application.
dimethoate	Do not apply within 28 days of grazing or harvesting for forage. Ground or aerial application.
malathion	Greatest efficacy at air temperatures above 20°C. Spray field margins of first-year clover in late summer or early fall when adult migration is underway. Remove cattle when treating; cattle may be returned immediately after treatment. Ground or aerial application.

References -

1. Craig, Pest. Res. Rep. 1971:179.
2. Craig, Pest. Res. Rep. 1964:136.
3. Craig, Pest. Res. Rep. 1965:135.
4. Swailes and McDonald, J. Econ. Entomol. 58:988, 1965.
5. Craig, Pest. Res. Rep. 1968: 150.
6. Bird, 80th Annual Rep., Entomol. Soc. Ont., 1949.
7. Craig, Can. Entomol. 110:883, 1978.

THRIPS IN RED CLOVER

Thrips in red clover seed fields

Economic Thresholds -

Damage is insignificant until 50-80 thrips/raceme are present. (1) Threshold levels occur only during years of early spring drought on dryland.

Reference -

1. Craig, unpublished data 1987, Agriculture and Agri-Food Canada, Saskatoon.